

Online Supplement

Detailed Results for Analyses in Paper

For paper “Sexual Orientation Differences in Treatment Expectation, Alliance, and Outcome.

Findings from Patients at Risk for Suicide in a Public Psychiatric Hospital”

Content

1	Comments to Statistical Analysis.....	3
2	Treatment Outcome – Difference Intake-Discharge Assessments.....	4
2.1	Change of Suicide Ideation.....	4
2.1.1	Interaction with Confounders.....	4
2.1.2	Adjusting for Confounders.....	5
2.2	Change of Hopelessness.....	7
2.2.1	Interaction with Confounders.....	7
2.2.2	Adjusting for Confounders.....	8
2.3	Change of Depression.....	10
2.3.1	Interaction with Confounders.....	10
2.3.2	Adjusting for Confounders.....	11
3	Responder Analysis.....	13
3.1	Suicide Ideation.....	13
3.1.1	Interactions with Confounders.....	14
3.1.2	Adjusting for Confounders.....	16
3.2	Hopelessness.....	18
3.2.1	Interactions with Confounders.....	18
3.2.2	Adjusting for Confounders.....	19
3.3	Depression.....	21
3.3.1	Interactions with Confounders.....	21
3.3.2	Adjusting for Confounders.....	22
4	Treatment Expectancy.....	24
3.4	Interaction with Confounders.....	24
3.4.1	Nationality.....	24
3.5	Adjusting for Confounders.....	25
5	Working Alliance.....	26
3.6	Interaction with Confounders.....	26
3.7	Adjusting for Confounders.....	28

1 Comments to Statistical Analysis

These supplemental statistical results are carried out with R 3.1.3 using additional packages such as “psych” and “rms” (for some descriptive tabulation), ggplot2 (for interaction plots), and “foreign” (for importing SPSS files).

2 Treatment Outcome – Difference Intake-Discharge Assessments

2.1 Change of Suicide Ideation

Linear Regression – SM Status as only predictor

Residuals:

Min	1Q	Median	3Q	Max
-28.797	-4.797	-3.305	5.203	27.203

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.7968	0.3475	13.804	<2e-16 ***
smyes	0.5085	0.7639	0.666	0.506

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 7.786 on 631 degrees of freedom

Multiple R-squared: 0.0007019, Adjusted R-squared: -0.0008818

F-statistic: 0.4432 on 1 and 631 DF, p-value: 0.5058

2.1.1 Interaction with Confounders

F7 Diagnosis

lm(formula = d_BSI ~ sm * F7)

Residuals:

Min	1Q	Median	3Q	Max
-28.83	-4.83	-3.00	5.17	27.17

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.8298	0.3559	13.569	<2e-16 ***
smyes	0.1702	0.7780	0.219	0.827
F7yes	-0.6375	1.5640	-0.408	0.684
smyes:F7yes	8.6375	3.8711	2.231	0.026 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 7.766 on 629 degrees of freedom

Multiple R-squared: 0.009005, Adjusted R-squared: 0.004278

F-statistic: 1.905 on 3 and 629 DF, p-value: 0.1275



2.1.2 Adjusting for Confounders

Full Multivariate Model

Call:

```
lm(formula = d_BSI ~ nationality + F3 + F7*sm + log(length_stay) * sm)
```

Residuals:

Min	1Q	Median	3Q	Max
-26.363	-5.141	-2.676	4.662	30.568

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.2972	2.1011	1.569	0.11710
nationality	-1.6587	0.9499	-1.746	0.08127 .
F3yes	2.1572	0.7224	2.986	0.00293 **
F7yes	-0.5045	1.5524	-0.325	0.74528
smyes	2.2559	3.6490	0.618	0.53666
log(length_stay)	0.5845	0.5730	1.020	0.30806
F7yes:smyes	8.1731	3.8412	2.128	0.03375 *
smyes:log(length_stay)	-0.6770	1.1548	-0.586	0.55790

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 7.701 on 625 degrees of freedom

Multiple R-squared: 0.0317, Adjusted R-squared: 0.02086

F-statistic: 2.923 on 7 and 625 DF, p-value: 0.00512



Adjusted to:F7=no sm=no length.stay=21

2.2 Change of Hopelessness

Linear Regression – SM Status as only predictor

Residuals:

Min	1Q	Median	3Q	Max
-28.797	-4.797	-3.305	5.203	27.203

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.7968	0.3475	13.804	<2e-16 ***
smyes	0.5085	0.7639	0.666	0.506

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 7.786 on 631 degrees of freedom

Multiple R-squared: 0.0007019, Adjusted R-squared: -0.0008818

F-statistic: 0.4432 on 1 and 631 DF, p-value: 0.5058

2.2.1 Interaction with Confounders

F6 Diagnosis

```
lm(formula = d_BHS ~ sm * F6)
```

Residuals:

Min	1Q	Median	3Q	Max
-13.5024	-3.5024	-0.5024	2.7634	15.4976

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.5024	0.2378	14.726	<2e-16 ***
smyes	0.7341	0.5542	1.325	0.1858
F6yes	0.2420	0.5617	0.431	0.6667
smyes:F6yes	-2.1891	1.0860	-2.016	0.0442 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.828 on 629 degrees of freedom

Multiple R-squared: 0.00733, Adjusted R-squared: 0.002596

F-statistic: 1.548 on 3 and 629 DF, p-value: 0.2009



2.2.2 Adjusting for Confounders

Full Multivariate Model

Call:

```
lm(formula = d_BHS ~ income + m_languag + F0 + F3 + F8 +
    F6 * sm + log(length_stay))
```

Residuals:

Min	1Q	Median	3Q	Max
-13.0885	-3.3232	-0.1569	3.0194	16.0740

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	4.3594531	1.2343305	3.532	0.000443	***
income	0.0004242	0.0001659	2.556	0.010811	*
m_languag	-1.6542639	0.6420080	-2.577	0.010204	*
F0yes	1.6797839	1.0501397	1.600	0.110198	
F3yes	0.8831951	0.4548372	1.942	0.052614	.
F8yes	-2.0280383	1.1485324	-1.766	0.077925	.
F6yes	0.6082591	0.5630221	1.080	0.280406	
smyes	0.8180816	0.5514322	1.484	0.138432	
log(length_stay)	-0.1074547	0.3098608	-0.347	0.728871	
F6yes:smyes	-2.3064893	1.0792971	-2.137	0.032984	*

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.76 on 623 degrees of freedom

Multiple R-squared: 0.04391, Adjusted R-squared: 0.0301

F-statistic: 3.179 on 9 and 623 DF, p-value: 0.0009075



Adjusted to:F6=no sm=no

2.3 Change of Depression

Linear Regression – SM Status as only predictor

Residuals:

Min	1Q	Median	3Q	Max
-13.5458	-3.5458	-0.5458	2.4542	15.4542

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.5458	0.2159	16.423	<2e-16 ***
smyes	0.1259	0.4746	0.265	0.791

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 4.837 on 631 degrees of freedom

Multiple R-squared: 0.0001116, Adjusted R-squared: -0.001473

F-statistic: 0.07042 on 1 and 631 DF, p-value: 0.7908

2.3.1 Interaction with Confounders

Length of stay

lm(formula = d_BDI ~ sm * length_stay)

Residuals:

Min	1Q	Median	3Q	Max
-29.366	-7.309	-0.534	6.677	34.391

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	12.16639	0.67357	18.063	<2e-16 ***
smyes	2.67140	1.42736	1.872	0.0617 .
length_stay	0.01427	0.01957	0.729	0.4664
smyes:length_stay	-0.08287	0.03400	-2.437	0.0151 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 10.78 on 629 degrees of freedom

Multiple R-squared: 0.01051, Adjusted R-squared: 0.005794

F-statistic: 2.228 on 3 and 629 DF, p-value: 0.08382



SM vs. heterosexuals: Length of Stay < 15 days $d = -0.28$
 > 30 days $d = 0.34$

2.3.2 Adjusting for Confounders

Full Multivariate Model

```
Call:
lm(formula = d_BDI ~ income + m_languag + F3 + log(length_stay) *
    sm)
```

Residuals:


Min	1Q	Median	3Q	Max
-30.992	-7.020	-0.964	6.579	33.256

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	14.1500874	2.9502233	4.796	2.02e-06 ***
income	0.0007955	0.0003645	2.182	0.02946 *
m_languag	-4.5701916	1.4198232	-3.219	0.00135 **
F3yes	3.9406123	0.9907661	3.977	7.78e-05 ***
log(length_stay)	-0.2130309	0.7837679	-0.272	0.78586
smyes	11.7819455	4.9793312	2.366	0.01828 *
log(length_stay):smyes	-3.6224037	1.5779521	-2.296	0.02203 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 10.54 on 626 degrees of freedom
 Multiple R-squared: 0.05895, Adjusted R-squared: 0.04993
 F-statistic: 6.536 on 6 and 626 DF, p-value: 1.046e-06



Adjusted to:length.stay=21 sm=no

3 Responder Analysis

Procedure according to Hiller, W., Schindler, A.C. & Lambert, M. J. (2011). Defining response and remission in psychotherapy research: A comparison of the RCI and the method of percent improvement. *Psychotherapy Research*, 22, 1 – 11.


Responder if:

1. baseline is in the pathological range and there is at least 50% improvement in the clinical range (baseline minus cut-off). For example, if a patient scores 30 on the BDI at baseline and 20 at follow-up (10-point difference), the cut-off is 14+, then this results in a percentual improvement of $10 / (30-14) * 100 = 62.5\%$
2. baseline is in the pathological range and the overall improvement has to be at least 25% in the general range (because if a patient's baseline is only slightly above the cut-off, then it is too easy to gain > 50% possible change in the clinical range). E.g, if a patient's baseline is 16 on the BDI and 14 after treatment, then this is an overall improvement of $(16-14)/16*100 = 12.5\%$ (but it would be 100% with criterion 1).
3. Patients can only be responders or nonresponders if the baseline is in the pathological range.

3.1 Suicide Ideation

Cutoff > 4 (arbitrary)

		res_BSI		
sm		1. responder	2. nonresponder	3. nonpathological
no		194	74	234
yes		57	26	48



Pearson's Chi-squared test

data: sm and res_BSI

X-squared = 4.6542, df = 2, p-value = 0.09758

3.1.1 Interactions with Confounders

F3 Diagnosis

```
lrm(formula = res_BSI_bin ~ sm * F3)
```

	Obs	351	Model Likelihood		Discrimination		Rank Discrim.	
			Ratio Test		Indexes		Indexes	
			LR chi2	9.51	R2	0.038	C	0.563
1. responder	251		d.f.	3	g	0.236	Dxy	0.126
2. nonresponder	100		Pr(> chi2)	0.0232	gr	1.267	gamma	0.221
max deriv	2e-11				gp	0.052	tau-a	0.052
					Brier	0.198		

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	-0.8473	0.3086	-2.75	0.0060
sm=yes	1.2993	0.5736	2.27	0.0235
F3=yes	-0.1441	0.3442	-0.42	0.6755
sm=yes * F3=yes	-1.5119	0.6625	-2.28	0.0225

Oddsratio without F3: 3.56 (1.16-11.67) (SM more likely nonresp.)
 Oddsratio with F3: 0.81 (0.41-1.53)

F6 Diagnosis

Logistic Regression Model

```
lrm(formula = res_BSI_bin ~ sm * F6)
```

Frequencies of Missing Values Due to Each Variable

	Obs	351	Model Likelihood		Discrimination		Rank Discrim.	
			Ratio Test		Indexes		Indexes	
			LR chi2	12.59	R2	0.051	C	0.594
1. responder	251		d.f.	3	g	0.366	Dxy	0.187
2. nonresponder	100		Pr(> chi2)	0.0056	gr	1.442	gamma	0.301
max deriv	1e-12				gp	0.077	tau-a	0.077
					Brier	0.196		

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	-1.0269	0.1601	-6.42	<0.0001
sm=yes	-0.4547	0.3852	-1.18	0.2378
F6=yes	0.2428	0.3082	0.79	0.4308
sm=yes * F6=yes	1.4464	0.5976	2.42	0.0155


Oddsratio without F6: 0.64 (0.29-1.33) (SM less likely nonresp.)
 Oddsratio with F6: 2.65 (1.08-1.68)

Length of Stay

```
lrm(formula = res_BSI_bin ~ sm * length_stay)
```

		Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs	351	LR chi2 10.76	R2 0.043	C 0.564
1. responder	251	d.f. 3	g 0.224	Dxy 0.128
2. nonresponder	100	Pr(> chi2) 0.0131	gr 1.252	gamma 0.145
max deriv	3e-04		gp 0.042	tau-a 0.052
			Brier 0.197	

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	-0.9977	0.1787	-5.58	<0.0001
sm=yes	-0.6452	0.4533	-1.42	0.1546
length_stay	0.0012	0.0042	0.30	0.7662
sm=yes * length_stay	0.0241	0.0115	2.09	0.0370



Cohen d responders 0.04 vs. nonresponders 0.69

3.1.2 Adjusting for Confounders

Full model

```
lrm(formula = res_BSI_bin ~ geschlecht + m_languag + F3 * sm +
      F6 * sm + log(length_stay) * sm)
```

Frequencies of Missing Values Due to Each Variable


res_BSI_bin	geschlecht	m_languag	F3	sm	F6	
length_stay						
282	0	0	0	0	0	0

		Model Likelihood	Discrimination	Rank Discrim.			
		Ratio Test	Indexes	Indexes			
Obs	351	LR chi2	31.60	R2	0.123	C	0.675
1. responder	251	d.f.	9	g	0.738	Dxy	0.351
2. nonresponder	100	Pr(> chi2)	0.0002	gr	2.092	gamma	0.354
max deriv	3e-07			gp	0.138	tau-a	0.143
				Brier	0.185		

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	-3.5367	1.0410	-3.40	0.0007
geschlecht	0.5224	0.2637	1.98	0.0476
m_languag	0.7328	0.3605	2.03	0.0421
F3=yes	-0.0887	0.3639	-0.24	0.8074
sm=yes	-2.8614	1.8020	-1.59	0.1123
F6=yes	0.1745	0.3233	0.54	0.5893
length_stay	0.3065	0.2259	1.36	0.1749
F3=yes * sm=yes	-1.0060	0.7764	-1.30	0.1951
sm=yes * F6=yes	0.9439	0.6868	1.37	0.1694
sm=yes * length_stay	0.9976	0.5269	1.89	0.0583

Effects				Response : res_BSI_bin			
Factor	Low	High	Diff.	Effect	S.E.	Lower 0.95	Upper 0.95
geschlecht	1	2	1	0.522420	0.26368	0.0056113	1.039200
Odds Ratio	1	2	1	1.686100	NA	1.0056000	2.827000
m_languag	1	2	1	0.732840	0.36049	0.0263010	1.439400
Odds Ratio	1	2	1	2.081000	NA	1.0267000	4.218100
length_stay	14	28	14	0.212430	0.15659	-0.0944910	0.519350
Odds Ratio	14	28	14	1.236700	NA	0.9098400	1.680900
F3 - no:yes	2	1	NA	0.088744	0.36395	-0.6245800	0.802060
Odds Ratio	2	1	NA	1.092800	NA	0.5354900	2.230100
sm - yes:no	1	2	NA	-0.830190	0.44766	-1.7076000	0.047195
Odds Ratio	1	2	NA	0.435970	NA	0.1813000	1.048300
F6 - yes:no	1	2	NA	0.174530	0.32327	-0.4590600	0.808120
Odds Ratio	1	2	NA	1.190700	NA	0.6318800	2.243700


Adjusted to: F3=yes sm=no F6=no length_stay=21



Adjusted to: F3=yes sm=no F6=no length.stay=21

3.2 Hopelessness

res_BHS		sm		
	1. responder	2. nonresponder	3. nonpathological	
no	93	290		119
yes	22	86		23



Pearson's Chi-squared test
 data: sm and res_BHS
 X-squared = 3.008, df = 2, p-value = 0.2222

3.2.1 Interactions with Confounders

F4 Diagnosis

```
lrm(formula = res_BHS_bin ~ sm * F4)
```

	Obs		Model Likelihood Ratio Test		Discrimination Indexes		Rank Discrim. Indexes	
			LR chi2	7.95	R2	0.024	C	0.568
1. responder	115	d.f.	3	g	0.292	Dxy	0.136	
2. nonresponder	376	Pr(> chi2)	0.0470	gr	1.340	gamma	0.222	
max deriv	4e-10			gp	0.049	tau-a	0.049	
				Brier	0.177			

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	1.1036	0.1410	7.83	<0.0001
sm=yes	0.7682	0.3678	2.09	0.0367
F4=yes	0.1157	0.2640	0.44	0.6613
sm=yes * F4=yes	-1.4279	0.5622	-2.54	0.0111

Oddsratio without F4: 2.12 (1.07-4.64) (SM more likely nonresp.)
 Oddsratio with F4: 0.51 (0.23-1.22)

3.2.2 Adjusting for Confounders

Full model

Logistic Regression Model

```
lrm(formula = res_BHS_bin ~ nationality + F0 + F4 * sm + log(length_stay))
```

Frequencies of Missing Values Due to Each Variable


res_BHS_bin	nationality	F0	F4	sm	length_stay
142	0	0	0	0	0

	Model Likelihood Ratio Test	Discrimination Indexes	Rank Discrim. Indexes
Obs	491	LR chi2 17.35	R2 0.052
1. responder	115	d.f. 6	g 0.499
2. nonresponder	376	Pr(> chi2) 0.0081	gr 1.647
max deriv	5e-09		gp 0.086
		Brier 0.173	tau-a 0.076

	Coef	S.E.	wald Z	Pr(> Z)
Intercept	-0.4853	0.7244	-0.67	0.5029
nationality	0.5947	0.3874	1.54	0.1248
F0=yes	-1.1124	0.5380	-2.07	0.0387
F4=yes	0.1624	0.2688	0.60	0.5456
sm=yes	0.6983	0.3710	1.88	0.0598
length_stay	0.3213	0.1856	1.73	0.0834
F4=yes * sm=yes	-1.4914	0.5672	-2.63	0.0086

Effects	Response : res_BHS_bin						
Factor	Low	High	Diff.	Effect	S.E.	Lower 0.95	Upper 0.95
nationality	1	2	1	0.59466	0.38738	-0.164580	1.353900
Odds Ratio	1	2	1	1.81240	NA	0.848250	3.872500
length_stay	14	28	14	0.22269	0.12862	-0.029413	0.474780
Odds Ratio	14	28	14	1.24940	NA	0.971010	1.607700
F0 - yes:no	1	2	NA	-1.11240	0.53798	-2.166800	-0.057957
Odds Ratio	1	2	NA	0.32877	NA	0.114540	0.943690
F4 - yes:no	1	2	NA	0.16244	0.26877	-0.364340	0.689220
Odds Ratio	1	2	NA	1.17640	NA	0.694650	1.992200
sm - yes:no	1	2	NA	0.69832	0.37100	-0.028819	1.425500
Odds Ratio	1	2	NA	2.01040	NA	0.971590	4.159800

Adjusted to: F4=no sm=no




Adjusted to:F3=yes sm=no F6=no length.stay=21

3.3 Depression

		res_BDI		
sm		1. responder	2. nonresponder	3.nonpathological
no		304	127	71
yes		84	37	10

Responder Analysis - Depression



Pearson's Chi-squared test

data: sm and res_BDI
X-squared = 4.0038, df = 2, p-value = 0.1351

3.3.1 Interactions with Confounders


Length of Stay

```
lrm(formula = res_BDI_bin ~ sm * length_stay)
Frequencies of Missing Values Due to Each Variable
```

res_BDI_bin	sm	length_stay
81	0	0

		Model Likelihood Ratio Test		Discrimination Indexes		Rank Discrim. Indexes	
Obs	552	LR chi2	15.34	R2	0.039	C	0.584
1. responder	388	d.f.	3	g	0.240	Dxy	0.168
2. nonresponder	164	Pr(> chi2)	0.0015	gr	1.271	gamma	0.196
max deriv	3e-08			gp	0.043	tau-a	0.070
				Brier	0.203		

	Coef	S.E.	Wald Z	Pr(> Z)
Intercept	-0.9010	0.1448	-6.22	<0.0001
sm=yes	-0.8932	0.4087	-2.19	0.0288
length_stay	0.0011	0.0039	0.29	0.7748
sm=yes * length_stay	0.0315	0.0121	2.59	0.0095



Cohen d responders 0.06 vs. nonresponders 0.73

3.3.2 Adjusting for Confounders

Full model

```
lrm(formula = res_BDI_bin ~ income + m_languag + F6 + log(length_stay)
*
sm)
```

Frequencies of Missing Values Due to Each Variable


res_BDI_bin	income	m_languag	F6	length_stay	sm
81	0	0	0	0	0

		Model Likelihood Ratio Test		Discrimination Indexes		Rank Discrim. Indexes	
Obs	552	LR chi2	52.36	R2	0.129	C	0.674
1. responder	388	d.f.	6	g	0.743	Dxy	0.348
2. nonresponder	164	Pr(> chi2)	<0.0001	gr	2.102	gamma	0.350
max deriv	0.02			gp	0.145	tau-a	0.145
				Brier	0.189		

	Coef	S.E.	wald z	Pr(> Z)
Intercept	-2.9233	0.7151	-4.09	<0.0001
income	-0.0002	0.0001	-1.84	0.0663
m_languag	1.3052	0.2995	4.36	<0.0001
F6=yes	0.4994	0.2312	2.16	0.0308
length_stay	0.2515	0.1838	1.37	0.1712
sm=yes	-3.7088	1.4048	-2.64	0.0083
length_stay * sm=yes	1.1305	0.4282	2.64	0.0083

Effects		Response : res_BDI_bin						
Factor	Low	High	Diff.	Effect	S.E.	Lower 0.95	Upper 0.95	
income	745.39	1537.2	791.82	-0.18265	0.099453	-0.377580	0.012271	
Odds Ratio	745.39	1537.2	791.82	0.83306		NA	0.685520 1.012300	
m_languag	1.00	2.0	1.00	1.30520	0.299500	0.718160	1.892200	
Odds Ratio	1.00	2.0	1.00	3.68830		NA	2.050700 6.633900	
length_stay	14.00	28.0	14.00	0.17435	0.127420	-0.075388	0.424090	
Odds Ratio	14.00	28.0	14.00	1.19050		NA	0.927380 1.528200	
F6 - yes:no	1.00	2.0	NA	0.49936	0.231230	0.046166	0.952550	
Odds Ratio	1.00	2.0	NA	1.64770		NA	1.047200 2.592300	
sm - yes:no	1.00	2.0	NA	-0.26707	0.262130	-0.780840	0.246700	
Odds Ratio	1.00	2.0	NA	0.76562		NA	0.458020 1.279800	

Adjusted to: length_stay=21 sm=no



Adjusted to: F3=yes sm=no F6=no length.stay=21

4 Treatment Expectancy

Linear Regression – SM Status as only predictor

Residuals:

Min	1Q	Median	3Q	Max
-73.127	-15.127	3.873	19.873	26.873

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	73.1275	1.0372	70.503	<2e-16 ***
smyes	0.5748	2.2800	0.252	0.801

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 23.24 on 631 degrees of freedom
Multiple R-squared: 0.0001007, Adjusted R-squared: -0.001484
F-statistic: 0.06356 on 1 and 631 DF, p-value: 0.801

3.4 Interaction with Confounders

3.4.1 Nationality

```
lm(formula = EXPECTANCY ~ sm * nationality)
```

Residuals:


Min	1Q	Median	3Q	Max
-72.778	-14.778	4.017	20.017	34.200

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	69.857	3.721	18.773	<2e-16 ***
smyes	16.309	8.265	1.973	0.0489 *
nationality	2.922	3.192	0.915	0.3604
smyes:nationality	-14.104	7.122	-1.980	0.0481 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 23.2 on 629 degrees of freedom
Multiple R-squared: 0.006298, Adjusted R-squared: 0.001558
F-statistic: 1.329 on 3 and 629 DF, p-value: 0.264



3.5 Adjusting for Confounders

Full Multivariate Model

```
lm(formula = EXPECTANCY ~ age + education + F1 + F3 + F6 + F8 +
    log(length.stay) + nationality * sm)
```

Residuals:

```
   Min       1Q   Median       3Q      Max
-76.76 -13.13   3.90   16.91   38.52
```

Coefficients:


	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	73.33841	6.63933	11.046	< 2e-16	***
age	0.33529	0.07592	4.416	1.19e-05	***
education	-2.71690	0.82170	-3.306	0.000999	***
F1yes	3.11889	2.02283	1.542	0.123619	
F3yes	-6.01239	2.17746	-2.761	0.005929	**
F6yes	-5.46526	2.33113	-2.344	0.019368	*
F8yes	-7.68336	5.39733	-1.424	0.155079	
log(length.stay)	-1.70657	1.46991	-1.161	0.246085	
nationality	2.66771	3.10681	0.859	0.390856	
smyes	16.52923	8.05765	2.051	0.040649	*
nationality:smyes	-12.69204	6.97115	-1.821	0.069140	.

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 22.52 on 622 degrees of freedom

Multiple R-squared: 0.07458, Adjusted R-squared: 0.0597

F-statistic: 5.013 on 10 and 622 DF, p-value: 4.996e-07



Adjusted to:nationality=1 sm=no

5 Working Alliance

Linear Regression – SM Status as only predictor

Residuals:

Min	1Q	Median	3Q	Max
-29.542	-5.542	1.458	7.458	12.458

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	47.5420	0.4093	116.140	<2e-16 ***
smyes	0.8301	0.9039	0.918	0.359

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 9.153 on 627 degrees of freedom

(4 observations deleted due to missingness)

Multiple R-squared: 0.001343, Adjusted R-squared: -0.0002495

F-statistic: 0.8433 on 1 and 627 DF, p-value: 0.3588

3.6 Interaction with Confounders

Nationality

```
lm(formula = WORKING_ALLIANCE ~ sm * nationality)
```

Residuals:

Min	1Q	Median	3Q	Max
-29.70	-5.70	1.30	7.30	16.27

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	49.017	1.460	33.583	<2e-16 ***
smyes	7.741	3.248	2.384	0.0174 *
nationality	-1.317	1.252	-1.052	0.2932
smyes:nationality	-6.196	2.794	-2.218	0.0269 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


Residual standard error: 9.094 on 625 degrees of freedom

(4 observations deleted due to missingness)

Multiple R-squared: 0.0173, Adjusted R-squared: 0.01259

F-statistic: 3.669 on 3 and 625 DF, p-value: 0.01217

F-statistic: 1.329 on 3 and 629 DF, p-value: 0.264



Length of stay

```
lm(formula = WORKING_ALLIANCE ~ sm * length_stay)
```

Residuals:

Min	1Q	Median	3Q	Max
-29.545	-5.756	1.175	7.217	14.620

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	48.18007	0.56586	85.144	<2e-16 ***
smyes	2.91048	1.20213	2.421	0.0158 *
length_stay	-0.02648	0.01643	-1.612	0.1075
smyes:length_stay	-0.06416	0.02856	-2.246	0.0250 *


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 9.041 on 625 degrees of freedom
(4 observations deleted due to missingness)

Multiple R-squared: 0.02876, Adjusted R-squared: 0.0241

F-statistic: 6.169 on 3 and 625 DF, p-value: 0.0003902

Interaction SM-Status x Length of Stay



3.7 Adjusting for Confounders

Full Multivariate Model

```
lm(formula = WORKING_ALLIANCE ~ age + income + mother.lang + F6 +
    nationality *
    sm + log(length.stay) * sm)
```

Residuals:

Min	1Q	Median	3Q	Max
-27.846	-5.006	1.324	6.783	15.705

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	53.1870257	2.8033334	18.973	< 2e-16	***
age	0.0372529	0.0300199	1.241	0.215098	
income	0.0008752	0.0003190	2.744	0.006251	**
mother.lang	-2.1360386	1.3435372	-1.590	0.112377	
F6yes	-3.2776815	0.9042098	-3.625	0.000313	***
nationality	-0.4411467	1.3462041	-0.328	0.743252	
smyes	12.7364565	5.0569053	2.519	0.012032	*
log(length.stay)	-1.6213674	0.6572585	-2.467	0.013900	*
nationality:smyes	-5.1899957	2.7390247	-1.895	0.058580	.
smyes:log(length.stay)	-1.6774997	1.3338478	-1.258	0.208996	


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 8.796 on 619 degrees of freedom

(4 observations deleted due to missingness)

Multiple R-squared: 0.08961, Adjusted R-squared: 0.07637

F-statistic: 6.77 on 9 and 619 DF, p-value: 2.678e-09



Adjusted to:nationality=1 sm=no length.stay=21